

兔抗 DOK2 (Phospho-Tyr299)多克隆抗体

中文名称：兔抗 DOK2 (Phospho-Tyr299)多克隆抗体

英文名称：Anti-DOK2 (Phospho-Tyr299) rabbit polyclonal antibody

别名：p56DOK; p56dok-2

相关类别：一抗

储存：冷冻（-20℃）避光

宿主：Rabbit

抗原：DOK2 (Phospho-Tyr299)

反应种属：Human

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

Background:	DOK proteins are enzymatically inert adaptor or scaffolding proteins. They provide a docking platform for the assembly of multimolecular signaling complexes. DOK2 may modulate the cellular proliferation induced by IL-4, as well as IL-2 and IL-3. May be involved in modulating Bcr-Abl signaling. Attenuates EGF-stimulated MAP kinase activation
Applications:	WB, IHC, IF
Name of antibody:	DOK2 (Phospho-Tyr299)
Immunogen:	Synthetic peptide of human DOK2 (Phospho-Tyr299)
Full name:	docking protein 2, 56kDa (Phospho-Tyr299)
Synonyms :	p56DOK; p56dok-2

SwissProt:	O60496
IHC positive control:	Human breast carcinoma
IHC Recommend dilution:	50-100
WB Predicted band size:	56 kDa
WB Positive control:	Jurkat cells treated with insulin
WB Recommended dilution:	500-1000
IF Positive control:	Hela cells
IF Recommended dilution	100-200



