

小鼠抗 MED28 单克隆抗体

中文名称： 小鼠抗 MED28 单克隆抗体

英文名称： Anti-MED28 mouse monoclonal antibody

别名： mediator complex subunit 28; EG1; magicin; 1500003D12Rik

相关类别： 一抗

储存： 冷冻（-20℃）

宿主： Mouse

抗原： MED28

反应种属： Human

标记物： Unconjugate

克隆类型： mouse monoclonal

技术规格

Background:

Component of the Mediator complex, a coactivator involved in the regulated transcription of nearly all RNA polymerase II-dependent genes. Mediator functions as a bridge to convey information from gene-specific regulatory proteins to the basal RNA polymerase II transcription machinery. Mediator is recruited to promoters by direct interactions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex with RNA polymerase II and the general transcription factors. May be part of a complex containing NF2/merlin that part

	icipates in cellular signaling to the actin cytoskeleton downstream of tyrosine kinase signaling pathways.
Applications:	WB, IHC
Name of antibody:	MED28
Immunogen:	Fusion protein of human MED28
Full name:	mediator complex subunit 28
Synonyms:	EG1; magicin; 1500003D12Rik
SwissProt:	Q9H204
IHC positive control:	Human colon tissue and Human colorectal cancer
IHC Recommend dilution:	50-150
WB Predicted band size:	20 KD
WB Positive control:	293T cell lysate
WB Recommended dilution:	500-2000