

订购热线: 4008-898-798

Anti-WDR18 antibody

Cat. No. ml223564

Package 25 μ l/100 μ l/200 μ l

Storage -20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Product overview

Description Anti-WDR18 rabbit polyclonal antibody

Applications ELISA, IHC

Immunogen Fusion protein of human WDR18

ReactivityHuman, MouseContent0.6 mg/mlHost speciesRabbit

Ig classImmunogen-specific rabbit IgGPurificationAntigen affinity purification

Target information

Symbol WDR18

Full name WD repeat domain 18

Synonyms Ipi3; R32184_1
Swissprot Q9BV38

Target Background

This gene encodes a member of the WD repeat protein family. WD repeats are minimally conserved regions of approximately 40 amino acids typically bracketed by gly-his and trp-asp (GH-WD), which may facilitate formation of heterotrimeric or multiprotein complexes. Members of this family are involved in a variety of cellular processes, including cell cycle progression, signal transduction, apoptosis, and gene regulation.




订购热线: 4008-898-798


Applications

Immunohistochemistry

Predicted cell location: Cytoplasm or Nucleus

Positive control: Human liver cancer Recommended dilution: 25-100





The image on the left is immunohistochemistry of paraffin-embedded Human liver cancer tissue using ml223564(WDR18 Antibody) at dilution 1/25, on the right is treated with fusion protein. (Original magnification: ×200)

ELISA

Recommended dilution: 2000-5000

联系电话: 4008-898-798, 021-61725725


联系QQ: 2881505695, 2881505696


邮箱: mlbio_cn@yeah.net 网址: www.mlbio.cn

Predicted cell location: Cytoplasm or Nucleus

Positive control: Human breast cancer

Recommended dilution: 25-100





The image on the left is immunohistochemistry of paraffin-embedded Human breast cancer tissue using ml223564(WDR18 Antibody) at dilution 1/25, on the right is treated with fusion protein. (Original

magnification: ×200)