

订购热线: 4008-898-798

Anti-CDC42EP2 antibody

Cat. No. ml224489

Package 25 μl/100 μl/200 μl

Storage -20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Product overview

Description Anti-CDC42EP2 rabbit polyclonal antibody

Applications ELISA, IHC

Immunogen Fusion protein of human CDC42EP2

Reactivity Human, Mouse, Rat

Content0.5 mg/mlHost speciesRabbit

Ig classImmunogen-specific rabbit IgGPurificationAntigen affinity purification

Target information

Symbol CDC42EP2

Full name CDC42 effector protein 2

Synonyms CEP2; BORG1
Swissprot O14613

Target Background

CDC42, a small Rho GTPase, regulates the formation of F-actin-containing structures through its interaction with the downstream effector proteins. The protein encoded by this gene is a member of the Borg family of CDC42 effector proteins. Borg family proteins contain a CRIB (Cdc42/Rac interactive-binding) domain. They bind to, and negatively regulate the function of CDC42. Coexpression of this protein with CDC42 suggested a role of this protein in actin filament assembly and cell shape control.





订购热线: 4008-898-798

Applications

Immunohistochemistry

Predicted cell location: Cytoplasm Positive control: Human liver cancer Recommended dilution: 25-100





The image on the left is immunohistochemistry of paraffin-embedded Human liver cancer tissue using ml224489(CDC42EP2 Antibody) at dilution 1/25, on the right is treated with fusion protein. (Original magnification: ×200)

ELISA

Recommended dilution: 5000-10000

联系电话: 4008-898-798, 021-61725725


联系QQ: 2881505695, 2881505696


邮箱: mlbio_cn@yeah.net 网址: www.mlbio.cn

Predicted cell location: Cytoplasm

Positive control: Human esophagus cancer

Recommended dilution: 25-100





The image on the left is immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using ml224489(CDC42EP2 Antibody) at dilution 1/25, on the right is treated with fusion protein. (Original magnification: ×200)