

订购热线: 4008-898-798

Anti-TAGAP antibody

Cat. No. ml224808

Package 25 μ l/100 μ l/200 μ l

Storage -20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Product overview

Description Anti-TAGAP rabbit polyclonal antibody

Applications ELISA, IHC

Immunogen Fusion protein of human TAGAP

ReactivityHuman, MouseContent0.54 mg/mlHost speciesRabbit

Ig classImmunogen-specific rabbit IgGPurificationAntigen affinity purification

Target information

Symbol TAGAP

Full name T-cell activation RhoGTPase activating protein
Synonyms FKSG15; IDDM21; TAGAP1; ARHGAP47

Swissprot Q8N103

Target Background

This gene encodes a member of the Rho GTPase-activator protein superfamily. The encoded protein may function as a Rho GTPase-activating protein. Alterations in this gene may be associated with several diseases, including rheumatoid arthritis, celiac disease, and multiple sclerosis. Alternate splicing results in multiple transcript variants encoding distinct isoforms.



订购热线: 4008-898-798

Applications


Immunohistochemistry

Predicted cell location: Cytoplasm and Cell membrane

Positive control: Human cervical cancer

Recommended dilution: 25-100





The image on the left is immunohistochemistry of paraffin-embedded Human cervical cancer tissue using ml224808(TAGAP Antibody) at dilution 1/20, on the right is treated with fusion protein. (Original magnification: ×200)

ELISA

Recommended dilution: 5000-10000


联系电话: 4008-898-798, 021-61725725


联系QQ: 2881505695, 2881505696

邮箱: mlbio_cn@yeah.net 网址: www.mlbio.cn

Predicted cell location: Cytoplasm and Cell membrane

Positive control: Human tonsil Recommended dilution: 25-100





The image on the left is immunohistochemistry of paraffin-embedded Human tonsil tissue using ml224808(TAGAP Antibody) at dilution 1/20, on the right is treated with fusion protein. (Original

magnification: ×200)