

## Anti-GCDH antibody

 Cat. No.
 ml224983

 Package
 25 μl/100 μl/200 μl

 Storage
 -20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Product overview Description Applications Immunogen Reactivity Content Host species Ig class Purification

Anti-GCDH rabbit polyclonal antibody ELISA, IHC Fusion protein of human GCDH Human, Mouse 1.02 mg/ml Rabbit Immunogen-specific rabbit IgG Antigen affinity purification

Target information Symbol Full name Synonyms Swissprot

GCDH glutaryl-CoA dehydrogenase GCD; ACAD5 Q92947

## **Target Background**


The protein encoded by this gene belongs to the acyl-CoA dehydrogenase family. It catalyzes the oxidative decarboxylation of glutaryl-CoA to crotonyl-CoA and CO(2) in the degradative pathway of L-lysine, L-hydroxylysine, and L-tryptophan metabolism. It uses electron transfer flavoprotein as its electron acceptor. The enzyme exists in the mitochondrial matrix as a homotetramer of 45-kD subunits. Mutations in this gene result in the metabolic disorder glutaric aciduria type 1, which is also known as glutaric acidemia type I. Alternative splicing of this gene results in multiple transcript variants. A related pseudogene has been identified on chromosome 12.



订购热线: 4008-898-798

Applications Immunohistochemistry Predicted cell location: Cytoplasm

Positive control: Human cervical cancer Recommended dilution: 30-150



The image on the left is immunohistochemistry of paraffin-embedded Human cervical cancer tissue using ml224983(GCDH Antibody) at dilution 1/30, on the right is treated with fusion protein. (Original magnification: ×200)

## ELISA

Recommended dilution: 5000-10000

联系电话: 4008-898-798, 021-61725725

联系QQ: 2881505695,2881505696

邮箱: mlbio\_cn@yeah.net 网址: www.mlbio.cn