

Anti-WNT10A antibody

 Cat. No.
 ml262611

 Package
 25 μl/100 μl/200 μl

 Storage
 -20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Target Background


The WNT gene family consists of structurally related genes which encode secreted signaling proteins. These proteins have been implicated in oncogenesis and in several developmental processes, including regulation of cell fate and patterning during embryogenesis. This gene is a member of the WNT gene family. It is strongly expressed in the cell lines of promyelocytic leukemia and Burkitt's lymphoma. In addition, it and another family member, the WNT6 gene, are strongly coexpressed in colorectal cancer cell lines. The gene overexpression may play key roles in carcinogenesis through activation of the WNT-beta-catenin-TCF signaling pathway. This gene and the WNT6 gene are clustered in the chromosome 2q35 region.



订购热线: 4008-898-798

Applications Immunohistochemistry

Predicted cell location: Cytoplasm Positive control: Human liver cancer Recommended dilution: 10-50



The image on the left is immunohistochemistry of paraffin-embedded Human liver cancer tissue using ml262611(WNT10A Antibody) at dilution 1/25, on the right is treated with synthetic peptide. (Original magnification: ×200)

ELISA

Recommended dilution: 1000-2000

联系电话: 4008-898-798, 021-61725725

联系QQ: 2881505695,2881505696

邮箱: mlbio_cn@yeah.net 网址: www.mlbio.cn