

订购热线: 4008-898-798

# Anti-DAZ4 antibody

**Cat. No.** ml120034

**Package** 25 μl/100 μl/200 μl

**Storage** -20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol

**Product overview** 

**Description** Anti-DAZ4 rabbit polyclonal antibody

Applications ELISA, WB, IHC

**Immunogen** Fusion protein of human DAZ4

Reactivity Human, Mouse

Content 1 mg/ml Host species Rabbit

Ig classImmunogen-specific rabbit IgGPurificationAntigen affinity purification

**Target information** 

Symbol DAZ4

Full name Deleted in azoospermia protein 4

**Synonyms** pDP1680, pDP1681

Swissprot Q86SG3

#### **Target Background**

This gene is a member of the DAZ gene family and is a candidate for the human Y-chromosomal azoospermia factor (AZF). Its expression is restricted to premeiotic germ cells, particularly in spermatogonia. It encodes an RNA-binding protein that is important for spermatogenesis. Four copies of this gene are found on chromosome Y within palindromic duplications; one pair of genes is part of the P2 palindrome and the second pair is part of the P1 palindrome. Each gene contains a 2.4 kb repeat including a 72-bp exon, called the DAZ repeat; the number of DAZ repeats is variable and there are several variations in the sequence of the DAZ repeat. Each copy of the gene also contains a 10.8 kb region that may be amplified; this region includes five exons that encode an RNA recognition motif (RRM) domain. This gene contains two copies of the 10.8 kb repeat. Alternatively spliced transcript variants encoding different isoforms have been described.





订购热线: 4008-898-798

#### **Applications**

## **Immunohistochemistry**

Predicted cell location: Cytoplasm Positive control: Human thyroid cancer Recommended dilution: 50-200





The image on the left is immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using ml120034(DAZ4 Antibody) at dilution 1/50, on the right is treated with fusion protein. (Original magnification: ×200)

## Western blotting

Predicted band size:65 kDa Positive control:TM4 cell lysate Recommended dilution: 500-2000

Gel: 8%SDS-PAGE


Lysate: 40 µg


Lane: TM4 cell lysate

elisakit producer Primary antibody: ml120034(DAZ4 Antibody) at dilution 1/200 Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution


Exposure time: 1 minute

Predicted cell location: Cytoplasm Positive control: Human liver cancer Recommended dilution: 50-200





The image on the left is immunohistochemistry of paraffin-embedded Human liver cancer tissue using ml120034(DAZ4 Antibody) at dilution 1/50, on the right is treated with fusion protein. (Original magnification: ×200)



#### **ELISA**

Recommended dilution: 5000-10000

联系电话: 4008-898-798, 021-61725725

联系QQ: 2881505695, 2881505696

邮箱: mlbio cn@yeah.net

网址: www.mlbio.cn