

订购热线: 4008-898-798

Anti-APPL1 antibody

Cat. No.	ml120280
Package	25 μl/100 μl/200 μl
Storage	-20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Product overview Description Applications Immunogen Reactivity Content **Host species** Ig class Purification

Anti-APPL1 rabbit polyclonal antibody ELISA, WB, IHC Fusion protein of human APPL1 Human, Mouse 0.1 mg/ml Rabbit Immunogen-specific rabbit IgG Antigen affinity purification

Target information	
Symbol	APPL1
Full name	Adaptor protein, phosphotyrosine interaction, PH domain and leucine zipper containing 1
Synonyms	APPL; DIP13alpha
Swissprot	Q9UKG1
Target Background	

Target Background

7000

The protein encoded by this gene has been shown to be involved in the regulation of cell proliferation, and in the crosstalk between the adiponectin signalling and insulin signalling pathways. The encoded protein binds many other proteins, including RAB5A, DCC, AKT2, PIK3CA, adiponectin receptors, and proteins of the NuRD/MeCP1 complex. This protein is found associated with endosomal membranes, but can be released by EGF and translocated to the nucleus.



订购热线: 4008-898-798

Applications

Immunohistochemistry Predicted cell location: Nucleus Positive control: Human esophagus cancer Recommended dilution: 100-300



The image on the left is immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using ml120280(APPL1 Antibody) at dilution 1/70, on the right is treated with fusion protein. (Original magnification: ×200)

Western blotting

Predicted band size:80 kDa Positive control:A172 cells Recommended dilution: 500-2000 Predicted cell location: Nucleus Positive control: Human lung cancer Recommended dilution: 100-300



kDa 250

130 95

72.

55



The image on the left is immunohistochemistry of paraffin-embedded Human lung cancer tissue using ml120280(APPL1 Antibody) at dilution 1/70, on the right is treated with fusion protein. (Original magnification: ×200)

Gel: 10%SDS-PAGE
Lysate: 40 µg
Lane: A172 cells
Primary antibody: ml120280(APPL1 Antibody) at dilution 1/600
Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution
Exposure time: 40 seconds
eisakit P
ELISA

ELISA Recommended dilution: 2000-5000

联系电话: 4008-898-798, 021-61725725

联系QQ: 2881505695, 2881505696

- 邮箱: mlbio_cn@yeah.net
- 网址: www.mlbio.cn