

# 订购热线: 4008-898-798

# Anti-CNN3 antibody

Cat. No.	ml121465
Package	25 μl/100 μl/200 μl
Storage	-20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Product overview Description Applications Immunogen Reactivity Content Host species Ig class Purification

Anti-CNN3 rabbit polyclonal antibody ELISA, WB, IHC Fusion protein of human CNN3 Human, Mouse, Rat 0.3 mg/ml Rabbit Immunogen-specific rabbit IgG Antigen affinity purification

Target information Symbol Full name Synonyms Swissprot

CNN3 Calponin 3, acidic

Q15417

### Target Background

This gene encodes a protein with a markedly acidic C terminus; the basic N-terminus is highly homologous to the N-terminus of a related gene, CNN1. Members of the CNN gene family all contain similar tandemly repeated motifs. This encoded protein is associated with the cytoskeleton but is not involved in contraction. Thin filament-associated protein that is implicated in the regulation and modulation of smooth muscle contraction. It is capable of binding to actin, calmodulin, troponin C and tropomyosin. The interaction of calponin with actin inhibits the actomyosin Mg-ATPase activity.




## 订购热线: 4008-898-798

### Applications

#### Immunohistochemistry

Predicted cell location: Cytoplasm Positive control: Human thyroid cancer Recommended dilution: 50-200



Predicted cell location: Cytoplasm Positive control: Human cervical cancer Recommended dilution: 50-200


The image on the left is immunohistochemistry of paraffin-embedded

Human cervical cancer tissue using ml121465(CNN3 Antibody) at


dilution 1/25, on the right is treated with fusion protein. (Original

2

1



magnification: ×200)



The image on the left is immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using ml121465(CNN3 Antibody) at dilution 1/25, on the right is treated with fusion protein. (Original magnification: ×200)

#### Western blotting

Predicted band size:36 kDa Positive control:Human fetal kidney and fetal muscle tissue Recommended dilution: 500-2000

> kDa 95 55-36 elisakit pri 28

Gel: 10%SDS-PAGE

Lysate: 40 µg

Lane 1-2: Human fetal kidney tissue, Human fetal muscle tissue Primary antibody: ml121465(CNN3 Antibody) at dilution 1/400 Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution Exposure time: 1 minute

**ELISA** 

Recommended dilution: 2000-5000

联系电话: 4008-898-798, 021-61725725

### 联系QQ: 2881505695, 2881505696

邮箱: mlbio cn@yeah.net

网址: www.mlbio.cn