

订购热线: 4008-898-798

# Anti-ALDH3A1 antibody

**Cat. No.** ml121567

Package 25 μl/100 μl/200 μl

**Storage** -20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol

**Product overview** 

**Description** Anti-ALDH3A1 rabbit polyclonal antibody

Applications ELISA, WB, IHC

Immunogen Fusion protein of human ALDH3A1

Reactivity Human, Mouse, Rat

Content0.2 mg/mlHost speciesRabbit

Ig classImmunogen-specific rabbit IgGPurificationAntigen affinity purification

**Target information** 

Symbol ALDH3A1

Full name aldehyde dehydrogenase 3 family, member A1

Synonyms ALDH3; ALDHIII

Swissprot P30838

#### **Target Background**

Aldehyde dehydrogenases oxidize various aldehydes to the corresponding acids. They are involved in the detoxification of alcohol-derived acetaldehyde and in the metabolism of corticosteroids, biogenic amines, neurotransmitters, and lipid peroxidation. The enzyme encoded by this gene forms a cytoplasmic homodimer that preferentially oxidizes aromatic and medium-chain (6 carbons or more) saturated and unsaturated aldehyde substrates. It is thought to promote resistance to UV and 4-hydroxy-2-nonenal-induced oxidative damage in the cornea. The gene is located within the Smith-Magenis syndrome region on chromosome 17. Multiple alternatively spliced variants, encoding the same protein, have been identified.




订购热线: 4008-898-798


## **Applications**

# **Immunohistochemistry**

Predicted cell location: Cytoplasm Positive control: Human thyroid cancer

Recommended dilution: 50-200





The image on the left is immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using ml121567(ALDH3A1 Antibody) at dilution 1/30, on the right is treated with fusion protein. (Original magnification: ×200)

## Western blotting

Predicted band size:50 kDa

Positive control:A549 and mouse eye tissue

Recommended dilution: 200-1000

Gel: 10%SDS-PAGE


Lysate: 40 µg

Lane 1-2: A549 cells, mouse eye tissue

Primary antibody: ml121567(ALDH3A1 Antibody) at dilution 1/350 uon d

Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution

Exposure time: 10 seconds



### **ELISA**

Recommended dilution: 2000-5000

联系电话: 4008-898-798, 021-61725725

联系QQ: 2881505695, 2881505696

邮箱: mlbio cn@yeah.net

网址: www.mlbio.cn