Anti-TICAM2 antibody

Cat. No.	ml122525
Package	25 µl/100 µl/200 µl
Storage	-20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Product overview

Description	Anti-TICAM2 rabbit polyclonal antibody
Applications	ELISA, WB, IHC
Immunogen	Fusion protein of human TICAM2
Reactivity	Human
Content	0.8 mg/ml
Host species	Rabbit
Ig class	Immunogen-specific rabbit IgG
Purification	Antigen affinity purification

Target information

Symbol	TICAM2
Full name	toll-like receptor adaptor molecule 2
Synonyms	TIRP; TRAM; TIRAP3; MyD88-4; TICAM-2
Swissprot	Q86XR7

Target Background

TIRP is a Toll/interleukin-1 receptor (IL1R; MIM 147810) (TIR) domain-containing adaptor protein involved in Toll receptor signaling

Applications

Immunohistochemistry

Predicted cell location: Cytoplasm

Positive control: Human thyroid cancer

Recommended dilution: 100-300

The image on the left is immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using ml122525(TICAM2 Antibody) at dilution 1/60, on the right is treated with fusion protein. (Original magnification: ×200)

Predicted cell location: Cytoplasm

Positive control: Human tonsil

Recommended dilution: 100-300

The image on the left is immunohistochemistry of paraffin-embedded Human tonsil tissue using ml122525(TICAM2 Antibody) at dilution 1/60, on the right is treated with fusion protein. (Original magnification: ×200)

Western blotting

Predicted band size:27 kDa

Positive control:Hela cells and human testis tissue, HT-29 and PC3 cells

Recommended dilution: 200-1000

Gel: 10%SDS-PAGE

Lysate: 40 μg

Lane 1-4: Hela cells, human testis tissue, HT29 cells, PC3 cells

Primary antibody: ml122525(TICAM2 Antibody) at dilution 1/200

Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution

Exposure time: 20 seconds

ELISA

Recommended dilution: 1000-2000

联系电话：**4008-898-798，021-61725725**

联系**QQ：2881505695，2881505696**

邮箱：**mlbio_cn@yeah.net**

网址：**www.mlbio.cn**