

订购热线: 4008-898-798

Anti-TPSB2 antibody

Cat. No. ml123395

Package 25 μl/100 μl/200 μl

Storage -20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Product overview

Description Anti-TPSB2 rabbit polyclonal antibody

Applications ELISA, WB, IHC

Immunogen Full length fusion protein

ReactivityHumanContent0.3 mg/mlHost speciesRabbit

Ig classImmunogen-specific rabbit IgGPurificationAntigen affinity purification

Target information

Symbol TPSB2

Full name tryptase beta 2 (gene/pseudogene)

Synonyms TPS2; tryptaseB; tryptaseC

Swissprot P20231

Target Background

Tryptases comprise a family of trypsin-like serine proteases, the peptidase family S1. Tryptases are enzymatically active only as heparin-stabilized tetramers, and they are resistant to all known endogenous proteinase inhibitors. Several tryptase genes are clustered on chromosome 16p13.3. These genes are characterized by several distinct features. They have a highly conserved 3' UTR and contain tandem repeat sequences at the 5' flank and 3' UTR which are thought to play a role in regulation of the mRNA stability. These genes have an intron immediately upstream of the initiator Met codon, which separates the site of transcription initiation from protein coding sequence. This feature is characteristic of tryptases but is unusual in other genes. The alleles of this gene exhibit an unusual amount of sequence variation, such that the alleles were once thought to represent two separate genes, beta II and beta III. Beta tryptases appear to be the main isoenzymes expressed in mast cells, whereas in basophils, alpha-tryptases predominate. Tryptases have been implicated as mediators in the pathogenesis of asthma and other allergic and inflammatory disorders.




订购热线: 4008-898-798


Applications

Immunohistochemistry

Predicted cell location: Secreted Positive control: Human thyroid cancer

Recommended dilution: 25-100





The image on the left is immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using ml123395(TPSB2 Antibody) at dilution 1/20, on the right is treated with fusion protein. (Original magnification: ×200)

Western blotting

Predicted band size:31 kDa

Positive control:Human left side lung cancer tissue lysate

Recommended dilution: 500-2000

Gel: 8%SDS-PAGE

Lysate: 20 µg

Lane: Human left side lung cancer tissue lysate

on disakil or Primary antibody: ml123395(TPSB2 Antibody) at dilution 1/200


Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution


Exposure time: 5 seconds

Predicted cell location: Secreted


Positive control: Human prostate cancer

Recommended dilution: 25-100





The image on the left is immunohistochemistry of paraffin-embedded Human prostate cancer tissue using ml123395(TPSB2 Antibody) at dilution 1/20, on the right is treated with fusion protein. (Original magnification: ×200)



ELISA

Recommended dilution: 2000-5000

联系电话: 4008-898-798, 021-61725725

联系QQ: 2881505695, 2881505696

邮箱: mlbio cn@yeah.net

网址: www.mlbio.cn