

# 订购热线: 4008-898-798

# Anti-ASB9 antibody

Cat. No.	ml123964
Package	25 μl/100 μl/200 μl
Storage	-20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Product overview
Description
Applications
Immunogen
Reactivity
Content
Host species
lg class
Purification

Anti-ASB9 rabbit polyclonal antibody ELISA, WB, IHC Full length fusion protein Human 0.5 mg/ml Rabbit Immunogen-specific rabbit IgG Antigen affinity purification

ducer

Target information	
Symbol	ASB9
Full name	ankyrin repeat and SOCS box containing 9
Synonyms	
Swissprot	Q96DX5

### **Target Background**

This gene encodes a member of the ankyrin repeat and suppressor of cytokine signaling (SOCS) box protein family. Members of this family can interact with the elongin B-C adapter complex via their SOCS box domain and further complex with the cullin and ring box proteins to form E3 ubiquitin ligase complexes. They may function to mediate the substraterecognition of the E3 ubiguitin ligases. A transcribed pseudogene of this gene has been identified on chromosome 15. Multiple transcript variants encoding different isoforms have been found for this gene.




## 订购热线: 4008-898-798

### Applications

Immunohistochemistry Predicted cell location: Cytoplasm Positive control: Human liver cancer

Recommended dilution: 25-100



The image on the left is immunohistochemistry of paraffin-embedded Human liver cancer tissue using ml123964(ASB9 Antibody) at dilution 1/25, on the right is treated with fusion protein. (Original magnification: ×200)

### Western blotting

Predicted band size:32 kDa Positive control:A549, A172 and Hepg2 cell Recor

ecommended dilution: 200-1000						
		kDa 95 —	1	2	3	
Gel: 12%SDS-PAGE		72 — 55 —				
Lysate: 40 µg		36 —				
Lane 1-3: A549, A172 and Hepg2 cell			_	-		
Primary antibody: ml123964(ASB9 Antibody) at dilution	on 1/400	20				
Secondary antibody: Goat anti rabbit IgG at 1/8000 d	lilution	17—				
Exposure time: 30 seconds	d'or					
		10—				
Lysate: 40 µg Lane 1-3: A549, A172 and Hepg2 cell Primary antibody: ml123964(ASB9 Antibody) at dilution Secondary antibody: Goat anti rabbit IgG at 1/8000 d		$\frac{72}{55}$ $36$ $28$ $17$	-		_	

ELISA

Recommended dilution: 5000-10000

### 联系电话: 4008-898-798, 021-61725725

## 联系QQ: 2881505695, 2881505696

邮箱: mlbio\_cn@yeah.net

网址: www.mlbio.cn