

订购热线: 4008-898-798

Anti-OCIAD1 antibody

Cat. No. ml122801

Package 25 μl/100 μl/200 μl

Storage -20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Product overview

Description Anti-OCIAD1 rabbit polyclonal antibody

Applications ELISA, WB

Immunogen Fusion protein of human OCIAD1

Reactivity Human, Mouse, Rat

Content0.3 mg/mlHost speciesRabbit

Ig classImmunogen-specific rabbit IgGPurificationAntigen affinity purification

Target information

Symbol OCIAD1

Full name OCIA domain containing 1
Synonyms OCIA; ASRIJ; TPA018

Swissprot Q9NX40

Target Background

OCIAD1 (OCIA domain containing 1), also known as OCIA (ovarian carcinoma immunoreactive antigen), TPA018 or ASRIJ, is a 245 amino acid endosomal protein that contains one OCIA domain and belongs to the OCIAD1 family. OCIAD1 exists as two alternatively spliced isoforms (termed OCIAD1 isoform A and B) that localize to different tissues; isoform A localizes to prostate, brain, mammary gland, testis, placenta and ovary, whereas isoform B is restricted to the central nervous system. The gene enocoding OCIAD1 maps to human chromosome 4, which represents approximately 6% of the human genome, contains nearly 900 genes and is associated with Huntington's disease, Ellis-van Creveld syndrome, methylmalonic acidemia and polycystic kidney disease.



订购热线: 4008-898-798

Applications

Western blotting

Predicted band size:28 kDa

Positive control:Hela cells, Jurkat cells and human placenta tissue, human testis tissue and 231 cells

Recommended dilution: 500-2000

Gel: 10%SDS-PAGE

Lysate: 40 µg

Lane 1-5: Hela cells, Jurkat cells, human placenta tissue,


human testis tissue, 231 cells

Primary antibody: ml122801(OCIAD1 Antibody) at dilution

1/200

Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution

Exposure time: 2 minutes



ELISA

Recommended dilution:2000-5000

联系电话: 4008-898-798, 021-61725725

联系QQ: 2881505695, 2881505696

邮箱: mlbio_cn@yeah.net

网址: www.mlbio.cn