

## Anti-GNB2L1 antibody

<b>Cat. No.</b>	ml122964
<b>Package</b>	25 µl/100 µl/200 µl
<b>Storage</b>	-20°C, pH7.4 PBS, 0.05% NaN <sub>3</sub> , 40% Glycerol

### Product overview

<b>Description</b>	Anti-GNB2L1 rabbit polyclonal antibody
<b>Applications</b>	ELISA, WB
<b>Immunogen</b>	Fusion protein of human GNB2L1
<b>Reactivity</b>	Human, Mouse, Rat
<b>Content</b>	0.5 mg/ml
<b>Host species</b>	Rabbit
<b>Ig class</b>	Immunogen-specific rabbit IgG
<b>Purification</b>	Antigen affinity purification

### Target information

<b>Symbol</b>	GNB2L1
<b>Full name</b>	guanine nucleotide binding protein (G protein), beta polypeptide 2-like 1
<b>Synonyms</b>	H12.3; HLC-7; PIG21; RACK1; Gnb2-rs1
<b>Swissprot</b>	P63244

### Target Background

RACK1 (receptor for activated C kinase 1) is a 317 amino acid G protein b subunit-like protein that functions as a RACK and inhibits the activity of Src tyrosine kinases. In response to PKC activation, the intracellular localization of RACK1 and PKC bII changes, and RACK1 and PKC bII co-localize to the same sites. RACK1 is therefore thought to be a shuttling protein for PKC bII. A deficit in RACK1 may be associated with impaired PKC activation in the aging brain.

订购热线: 4008-898-798

#### Applications

##### Western blotting

Predicted band size: 35 kDa

Positive control: NIH/3T3 and Jurkat cells

Recommended dilution: 200-1000

Gel: 12%SDS-PAGE


Lysate: 50 µg

Lane 1-2: NIH/3T3 cells, Jurkat cells

Primary antibody: ml122964(GNB2L1 Antibody) at dilution 1/300

Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution

Exposure time: 20 seconds



#### ELISA

Recommended dilution: 1000-2000

联系电话: 4008-898-798, 021-61725725

联系QQ: 2881505695, 2881505696

邮箱: [mlbio\\_cn@yeah.net](mailto:mlbio_cn@yeah.net)

网址: [www.mlbio.cn](http://www.mlbio.cn)