

# 订购热线: 4008-898-798

# Anti-PDE4D antibody

Cat. No.	ml162030
Package	25 μl/100 μl/200 μl
Storage	-20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Product overview	
Description	
Applications	
Immunogen	
Reactivity	
Content	
Host species	
lg class	
Purification	

Anti-PDE4D rabbit polyclonal antibody ELISA, WB, IHC Synthetic peptide of human PDE4D Human 0.2 mg/ml Rabbit Immunogen-specific rabbit IgG Antigen affinity purification

Target information	
Symbol	PDE4D
Full name	phosphodiesterase 4D, cAMP-specific
Synonyms	DPDE3; PDE43; STRK1; ACRDYS2; HSPDE4D; PDE4DN2
Swissprot	Q08499
Target Background	duce

### **Target Background**

This gene encodes one of four mammalian counterparts to the fruit fly 'dunce' gene. The encoded protein has 3',5'-cyclic-AMP phosphodiesterase activity and degrades cAMP, which acts as a signal transduction molecule in multiple cell types. This gene uses different promoters to generate multiple alternatively spliced transcript variants that encode functional Goodei proteins.




## 订购热线: 4008-898-798


### Applications

Immunohistochemistry

Predicted cell location: Cytoplasm Positive control: Human prostate cancer Recommended dilution: 25-100



Predicted cell location: Cytoplasm Positive control: Human brain Recommended dilution: 25-100




130-

95-

72-

55-



The image on the left is immunohistochemistry of paraffin-embedded Human prostate cancer tissue using ml162030(PDE4D Antibody) at dilution 1/20, on the right is treated with synthetic peptide. (Original magnification: ×200)

#### Western blotting

Predicted band size:91 kDa Positive control:Hela cells Recommended dilution: 200-1000 The image on the left is immunohistochemistry of paraffin-embedded Human brain tissue using ml162030(PDE4D Antibody) at dilution 1/20, on the right is treated with synthetic peptide. (Original magnification: ×200)

Gel: 6%SDS-PAGE Lysate: 40 µg

Lane: Hela cells

in on other Primary antibody: ml162030(PDE4D Antibody) at dilution 1/200 Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution Exposure time: 1 minute

**ELISA** Recommended dilution: 1000-2000

## 联系电话: 4008-898-798, 021-61725725

### 联系QQ: 2881505695, 2881505696

邮箱: mlbio cn@yeah.net

网址: www.mlbio.cn