

Anti-CCDC47 antibody

Cat. No.	ml125084
Package	25 µl/100 µl/200 µl
Storage	-20°C, pH7.4 PBS, 0.05% NaN ₃ , 40% Glycerol

Product overview

Description	Anti-CCDC47 rabbit polyclonal antibody
Applications	ELISA, WB
Immunogen	Fusion protein of human CCDC47
Reactivity	Human, Mouse, Rat
Content	0.72 mg/ml
Host species	Rabbit
Ig class	Immunogen-specific rabbit IgG
Purification	Antigen affinity purification

Target information

Symbol	CCDC47
Full name	coiled-coil domain containing 47
Synonyms	GK001; MSTP041
Swissprot	Q96A33

Target Background

The coiled-coil domain is a common protein motif that is often involved in protein oligomerization and is found in proteins such as transcription factors and intermediate filaments. The CCDC47 gene maps to chromosome 17 at 17q23.3. Little is known about this single-pass membrane protein except that the coiled-coil domain is within the cytosolic domain near the carboxy terminus.

订购热线: 4008-898-798

Applications

Western blotting

Predicted band size: 56 kDa

Positive control: K562 and A172 cell, Human bladder transitional cell carcinoma grade 2-3 tissue, NIH/3T3 cell, Mouse liver tissue lysates

Recommended dilution: 500-2000

Gel: 8%SDS-PAGE


Lysate: 40 µg

Lane 1-5: K562 and A172 cell, Human bladder transitional cell carcinoma grade 2-3 tissue, NIH/3T3 cell, Mouse liver tissue lysates

Primary antibody: ml125084(CCDC47 Antibody) at dilution 1/300

Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution

Exposure time: 5 seconds



ELISA

Recommended dilution: 5000-10000

联系电话: 4008-898-798, 021-61725725

联系QQ: 2881505695, 2881505696

邮箱: mlbio_cn@yeah.net

网址: www.mlbio.cn

mlbio 酶联生物
Good elisakit producers
订购热线: 4008-898-798