

Anti-ACSF2 antibody

Cat. No.	ml125125
Package	25 µl/100 µl/200 µl
Storage	-20°C, pH7.4 PBS, 0.05% NaN ₃ , 40% Glycerol

Product overview

Description	Anti-ACSF2 rabbit polyclonal antibody
Applications	ELISA, WB, IHC
Immunogen	Fusion protein of human ACSF2
Reactivity	Human, Mouse, Rat
Content	0.96 mg/ml
Host species	Rabbit
Ig class	Immunogen-specific rabbit IgG
Purification	Antigen affinity purification

Target information

Symbol	ACSF2
Full name	acyl-CoA synthetase family member 2
Synonyms	ACSMW; AVYV493
Swissprot	Q96CM8

Target Background

Acyl-CoA synthases catalyze the initial reaction in fatty acid metabolism, by forming a thioester with CoA. Has some preference toward medium-chain substrates. Plays a role in adipocyte differentiation.

订购热线: 4008-898-798


Applications

Immunohistochemistry

Predicted cell location: Cytoplasm

Positive control: Human thyroid cancer

Recommended dilution: 30-150



The image on the left is immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using ml125125(ACSF2 Antibody) at dilution 1/30, on the right is treated with fusion protein. (Original magnification: ×200)

Western blotting

Predicted band size: 68 kDa

Positive control: Human thyroid tissue, 293T cell, Rat liver tissue, A172 and Jurkat cell lysates

Recommended dilution: 500-2000

Gel: 8%SDS-PAGE


Lysate: 40 µg

Lane 1-5: Human thyroid tissue, 293T cell, Rat liver tissue, A172 and Jurkat cell lysates

Primary antibody: ml125125(ACSF2 Antibody) at dilution 1/400

Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution

Exposure time: 30 seconds



ELISA

Recommended dilution: 5000-10000

联系电话: 4008-898-798, 021-61725725

联系QQ: 2881505695, 2881505696

邮箱: mlbio_cn@yeah.net

网址: www.mlbio.cn