

订购热线: 4008-898-798

# Anti-HLA-DRB4 antibody

**Cat. No.** ml125381

Package 25  $\mu$ l/100  $\mu$ l/200  $\mu$ l

**Storage** -20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol

**Product overview** 

**Description** Anti-HLA-DRB4 rabbit polyclonal antibody

**Applications** ELISA, WB

Immunogen Fusion protein of human HLA-DRB4

Reactivity Human

Content 0.84 mg/ml

Host species Rabbit

Ig classImmunogen-specific rabbit IgGPurificationAntigen affinity purification

**Target information** 

Symbol HLA-DRB4

Full name major histocompatibility complex, class II, DR beta 4

Synonyms DR4; DRB4; HLA-DR4B

Swissprot P13762

### **Target Background**

HLA-DRB4 belongs to the HLA class II beta chain paralogues. This class II molecule is a heterodimer consisting of an alpha (DRA) and a beta (DRB) chain, both anchored in the membrane. It plays a central role in the immune system by presenting peptides derived from extracellular proteins. Class II molecules are expressed in antigen presenting cells (APC: B lymphocytes, dendritic cells, macrophages). The beta chain is approximately 26-28 kDa and its gene contains 6 exons. Exon one encodes the leader peptide, exons 2 and 3 encode the two extracellular domains, exon 4 encodes the transmembrane domain and exon 5 encodes the cytoplasmic tail. Within the DR molecule the beta chain contains all the polymorphisms specifying the peptide binding specificities. Typing for these polymorphisms is routinely done for bone marrow and kidney transplantation. DRB1 is expressed at a level five times higher than its paralogues DRB3, DRB4 and DRB5. The presence of DRB4 is linked with allelic variants of DRB1, otherwise it is omitted. There are 4 related pseudogenes: DRB2, DRB6, DRB7, DRB8 and DRB9.



订购热线: 4008-898-798

## **Applications**

### Western blotting

Predicted band size:30 kDa
Positive control:Raji cell lysate
Recommended dilution: 500-2000

Gel: 8%SDS-PAGE

Lysate: 40 µg

Lane: Raji cell lysate

Primary antibody: ml125381(HLA-DRB4 Antibody) at

dilution 1/350

Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution

Exposure time: 5 seconds

kDa 95 — 72 — 55 — 36 — 28 — 17 —

# **ELISA**

Recommended dilution:5000-10000

联系电话: 4008-898-798, 021-61725725

联系QQ: 2881505695, 2881505696

邮箱: mlbio\_cn@yeah.net

网址: www.mlbio.cn