

# 订购热线: 4008-898-798

# Anti-DEFA1 antibody

Cat. No. ml160252 Package 25 µl/100 µl/200 µl Storage -20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol

**Product overview** Description Applications Immunogen Reactivity Content **Host species** Ig class Purification

Anti-DEFA1 rabbit polyclonal antibody ELISA, WB, IHC Synthetic peptide of human DEFA1 Human 0.4 mg/ml Rabbit Immunogen-specific rabbit IgG Antigen affinity purification

Target information	
Symbol	DEFA1
Full name	Defensin, alpha 1
Synonyms	HP1, MRS, DEF1, HP-1, DEFA2, HNP-
Swissprot	P59665
Target Background	ducers

#### **Target Background**


Defensins are a family of microbicidal and cytotoxic peptides thought to be involved in host defense. They are abundant in the granules of neutrophils and also found in the epithelia of mucosal surfaces such as those of the intestine, respiratory tract, urinary tract, and vagina. Members of the defensin family are highly similar in protein sequence and distinguished by a conserved cysteine motif. The protein encoded by this gene, defensin, alpha 1, is found in the microbicidal granules of neutrophils and likely plays a role in phagocyte-mediated host defense. Several alpha defensin genes are clustered on chromosome 8. This gene differs from defensin, alpha 3 by only one amino acid. This gene and the gene encoding defensin, alpha 3 are both subject to copy number variation.



## 订购热线: 4008-898-798

### Applications

Immunohistochemistry Predicted cell location: Cytoplasm Positive control: Human esophagus cancer Recommended dilution: 25-50



The image on the left is immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using ml160252(DEFA1 Antibody) at dilution 1/15, on the right is treated with synthetic peptide. (Original magnification: ×200)

### W

Vestern blotting	s der '
Predicted band size:10 kDa	
Positive control:Human lung tissue	
Recommended dilution: 500-1000	
	KDa
	40—
Gel: 12%SDS-PAGE	25
Lysate: 30 µg	35—
Lane: Human lung tissue	25—
Primary antibody: ml160252(DEFA1 Antibody) at dilution 1/550	
Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution	15—
Exposure time: 20 seconds	10
akit P.	1000

ELISA Recommended dilution: 500-5000

### 联系电话: 4008-898-798, 021-61725725

## 联系QQ: 2881505695, 2881505696

- 邮箱: mlbio\_cn@yeah.net
- 网址: www.mlbio.cn