

## 订购热线: 4008-898-798

# Anti-ESR2 antibody

Cat. No.	ml160315
Package	25 μl/100 μl/200 μl
Storage	-20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol

**Product overview** Description Applications Immunogen Reactivity Content **Host species** Ig class Purification

Anti-ESR2 rabbit polyclonal antibody ELISA, WB, IHC Synthetic peptide of human ESR2 Human, Mouse, Rat 0.1 mg/ml Rabbit Immunogen-specific rabbit IgG Antigen affinity purification

Target information	
Symbol	ESR2
Full name	estrogen receptor 2
Synonyms	Erb; ESRB; ESTRB; NR3A2; ER-BETA; ESR-BETA
Swissprot	Q92731
Target Background	duc

#### **Target Background**


This gene encodes a member of the family of estrogen receptors and superfamily of nuclear receptor transcription factors. The gene product contains an N-terminal DNA binding domain and C-terminal ligand binding domain and is localized to the nucleus, cytoplasm, and mitochondria. Upon binding to 17beta-estradiol or related ligands, the encoded protein forms homo- or hetero-dimers that interact with specific DNA sequences to activate transcription. Some isoforms dominantly inhibit the activity of other estrogen receptor family members. Several alternatively spliced transcript variants of this gene have been described, but the full-length nature of some of these variants has not been fully characterized.



## 订购热线: 4008-898-798

### Applications

Immunohistochemistry Predicted cell location: Nucleus Positive control: Human esophagus cancer Recommended dilution: 25-100




The image on the left is immunohistochemistry of paraffinembedded Human esophagus cancer tissue using ml160315(ESR2 Antibody) at dilution 1/20, on the right is treated with synthetic peptide. (Original magnification: ×200)

#### Western blotting

Predicted band size:59 kDa Positive control:A549,PC-3,K562 and 231 cell lysates Recommended dilution: 200-1000

Gel: 8%SDS-PAGE Lysate: 40 µg elisatit producer Lane 1-4: A549, PC-3, K562 and 231 cell lysates Primary antibody: ml160315(ESR2 Antibody) at dilution 1/300 Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution Exposure time: 1 minute



**ELISA** 

Recommended dilution: 5000-10000

### 联系电话: 4008-898-798, 021-61725725

### 联系QQ: 2881505695, 2881505696

邮箱: mlbio cn@yeah.net

网址: www.mlbio.cn