

Anti-ABCD4 antibody

Cat. No.	ml160807
Package	25 µl/100 µl/200 µl
Storage	-20°C, pH7.4 PBS, 0.05% NaN ₃ , 40% Glycerol

Product overview

Description	Anti-ABCD4 rabbit polyclonal antibody
Applications	ELISA, WB
Immunogen	Synthetic peptide of human ABCD4
Reactivity	Human, Mouse
Content	0.3 mg/ml
Host species	Rabbit
Ig class	Immunogen-specific rabbit IgG
Purification	Antigen affinity purification

Target information

Symbol	ABCD4
Full name	ATP-binding cassette, sub-family D (ALD), member 4
Synonyms	P70R; P79R; ABC41; MAHCJ; PMP69; PXMP1L; EST352188
Swissprot	O14678

Target Background

The protein encoded by this gene is a member of the superfamily of ATP-binding cassette (ABC) transporters. ABC proteins transport various molecules across extra- and intra-cellular membranes. ABC genes are divided into seven distinct subfamilies (ABC1, MDR/TAP, MRP, ALD, OABP, GCN20, White). This protein is a member of the ALD subfamily, which is involved in peroxisomal import of fatty acids and/or fatty acyl-CoAs in the organelle. All known peroxisomal ABC transporters are half transporters which require a partner half transporter molecule to form a functional homodimeric or heterodimeric transporter. The function of this peroxisomal membrane protein is unknown. However, it is speculated that it may function as a heterodimer for another peroxisomal ABC transporter and, therefore, may modify the adrenoleukodystrophy phenotype. It may also play a role in the process of peroxisome biogenesis.

订购热线: 4008-898-798

Applications

Western blotting

Predicted band size:69 kDa

Positive control:Jurkat cells

Recommended dilution: 500-2000

Gel: 10%SDS-PAGE


Lysate: 40 μ g

Lane: Jurkat cells

Primary antibody: ml160807(ABCD4 Antibody) at dilution 1/700

Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution

Exposure time: 1 minute



ELISA

Recommended dilution:2000-5000

联系电话: 4008-898-798, 021-61725725

联系QQ: 2881505695, 2881505696

邮箱: mlbio_cn@yeah.net

网址: www.mlbio.cn