

Anti-DDIT4 antibody

Cat. No.	ml122220
Package	25 µl/100 µl/200 µl
Storage	-20°C, pH7.4 PBS, 0.05% NaN ₃ , 40% Glycerol

Product overview

Description	Anti-DDIT4 rabbit polyclonal antibody
Applications	ELISA, IHC
Immunogen	Fusion protein of human DDIT4
Reactivity	Human, Mouse, Rat
Content	0.8 mg/ml
Host species	Rabbit
Ig class	Immunogen-specific rabbit IgG
Purification	Antigen affinity purification

Target information

Symbol	DDIT4
Full name	DNA-damage-inducible transcript 4
Synonyms	Dig2; REDD1; REDD-1
Swissprot	Q9NX09

Target Background

REDD-1, also designated DNA-damage-inducible transcript 4, dig2 or RTP801, is thought to function in the regulation of reactive oxygen species (ROS). REDD-1 expression has also been linked to apoptosis, Ab toxicity and the pathogenesis of ischemic diseases. As an HIF-1-responsive gene, REDD-1 exhibits strong hypoxia-dependent upregulation in ischemic cells of neuronal origin. In response to stress due to DNA damage and glucocorticoid treatment, REDD-1 is upregulated at the transcriptional level. REDD-1 negatively regulates the mammalian target of Rapamycin (mTOR), a serine/threonine kinase often referred to as FRAP. It is crucial in the coupling of extra- and intracellular cues to FRAP regulation. The absence of REDD-1 is associated with the development of retinopathy, a major cause of blindness.



Applications

Immunohistochemistry

Predicted cell location: Cytoplasm

Positive control: Human liver cancer

Recommended dilution: 50-200





The image on the left is immunohistochemistry of paraffin-embedded Human liver cancer tissue using ml122220(DDIT4 Antibody) at dilution 1/30, on the right is treated with fusion protein. (Original magnification: $\times 200$)

Predicted cell location: Cytoplasm

Positive control: Human gastric cancer

Recommended dilution: 50-200



The image on the left is immunohistochemistry of paraffin-embedded Human gastric cancer tissue using ml122220(DDIT4 Antibody) at dilution 1/30, on the right is treated with fusion protein. (Original magnification: $\times 200$)

ELISA

Recommended dilution: 2000-5000

联系电话: 4008-898-798, 021-61725725

联系QQ: 2881505695, 2881505696

邮箱: mlbio_cn@yeah.net

网址: www.mlbio.cn