

Anti-CASP14 antibody

| | |
|-----------------|---|
| Cat. No. | ml220081 |
| Package | 25 µl/100 µl/200 µl |
| Storage | -20°C, pH7.4 PBS, 0.05% NaN ₃ , 40% Glycerol |

Product overview

| | |
|---------------------|--|
| Description | Anti-CASP14 rabbit polyclonal antibody |
| Applications | ELISA, WB |
| Immunogen | Fusion protein of human CASP14 |
| Reactivity | Human |
| Content | 0.2 mg/ml |
| Host species | Rabbit |
| Ig class | Immunogen-specific rabbit IgG |
| Purification | Antigen affinity purification |

Target information

| | |
|------------------|--|
| Symbol | CASP14 |
| Full name | caspase 14, apoptosis-related cysteine peptidase |
| Synonyms | |
| Swissprot | P31944 |

Target Background

This gene encodes a member of the cysteine-aspartic acid protease (caspase) family. Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes which undergo proteolytic processing at conserved aspartic residues to produce two subunits, large and small, that dimerize to form the active enzyme. This caspase has been shown to be processed and activated by caspase 8 and caspase 10 in vitro, and by anti-Fas agonist antibody or TNF-related apoptosis inducing ligand in vivo. The expression and processing of this caspase may be involved in keratinocyte terminal differentiation, which is important for the formation of the skin barrier.

订购热线: 4008-898-798

Applications

Western blotting

Predicted band size: 28 kDa

Positive control: Jurkat, A431 and A375 cells

Recommended dilution: 200-1000

Gel: 12% SDS-PAGE


Lysate: 40 µg

Lane 1-3: Jurkat cells, A431 cells, A375 cells

Primary antibody: ml220081(CASP14 Antibody) at dilution 1/300

Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution

Exposure time: 20 seconds



ELISA

Recommended dilution: 1000-2000

联系电话: 4008-898-798, 021-61725725

联系QQ: 2881505695, 2881505696

邮箱: mlbio_cn@yeah.net

网址: www.mlbio.cn