

## Anti-TFAP2C antibody

<b>Cat. No.</b>	ml161286
<b>Package</b>	25 µl/100 µl/200 µl
<b>Storage</b>	-20°C, pH7.4 PBS, 0.05% NaN <sub>3</sub> , 40% Glycerol

### Product overview

<b>Description</b>	Anti-TFAP2C rabbit polyclonal antibody
<b>Applications</b>	ELISA, WB
<b>Immunogen</b>	Synthetic peptide of human TFAP2C
<b>Reactivity</b>	Human, Mouse
<b>Content</b>	0.583 mg/ml
<b>Host species</b>	Rabbit
<b>Ig class</b>	Immunogen-specific rabbit IgG
<b>Purification</b>	Antigen affinity purification

### Target information

<b>Symbol</b>	TFAP2C
<b>Full name</b>	transcription factor AP-2 gamma (activating enhancer binding protein 2 gamma)
<b>Synonyms</b>	ERF1; TFAP2G; hAP-2g; AP2-GAMMA
<b>Swissprot</b>	Q92754

### Target Background

The protein encoded by this gene is a sequence-specific DNA-binding transcription factor involved in the activation of several developmental genes. The encoded protein can act as either a homodimer or heterodimer with other family members and is induced during retinoic acid-mediated differentiation. It plays a role in the development of the eyes, face, body wall, limbs, and neural tube.

订购热线: 4008-898-798

### Applications

#### Western blotting

Predicted band size:49 kDa

Positive control:Human fetal muscle tissue, huvec and hela cells

Recommended dilution: 200-1000

Gel: 10%SDS-PAGE


Lysate: 40 µg

Lane 1-3: Human fetal muscle tissue, huvec cells, hela cells

Primary antibody: ml161286(TFAP2C Antibody) at dilution 1/270

Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution

Exposure time: 30 seconds



#### ELISA

Recommended dilution:1000-2000

联系电话: 4008-898-798, 021-61725725

联系QQ: 2881505695, 2881505696

邮箱: [mlbio\\_cn@yeah.net](mailto:mlbio_cn@yeah.net)

网址: [www.mlbio.cn](http://www.mlbio.cn)