

Anti-EHHADH antibody

Cat. No.	ml222276
Package	25 µl/100 µl/200 µl
Storage	-20°C, pH7.4 PBS, 0.05% NaN ₃ , 40% Glycerol

Product overview

Description	Anti-EHHADH rabbit polyclonal antibody
Applications	ELISA, WB, IHC
Immunogen	Fusion protein of human EHHADH
Reactivity	Human, Mouse
Content	0.5 mg/ml
Host species	Rabbit
Ig class	Immunogen-specific rabbit IgG
Purification	Antigen affinity purification

Target information

Symbol	EHHADH
Full name	enoyl-CoA, hydratase/3-hydroxyacyl CoA dehydrogenase

Synonyms LBP; ECHD; LBFP; PBFE; L-PBE

Swissprot Q08426

Target Background

The protein encoded by this gene is a bifunctional enzyme and is one of the four enzymes of the peroxisomal beta-oxidation pathway. The N-terminal region of the encoded protein contains enoyl-CoA hydratase activity while the C-terminal region contains 3-hydroxyacyl-CoA dehydrogenase activity. Defects in this gene are a cause of peroxisomal disorders such as Zellweger syndrome. Two transcript variants encoding different isoforms have been found for this gene.


Applications

Immunohistochemistry

Predicted cell location: Cytoplasm

Positive control: Human liver cancer

Recommended dilution: 50-200




The image on the left is immunohistochemistry of paraffin-embedded Human liver cancer tissue using ml222276(EHHADH Antibody) at dilution 1/20, on the right is treated with fusion protein. (Original magnification: $\times 200$)

Predicted cell location: Cytoplasm

Positive control: Human thyroid cancer

Recommended dilution: 50-200



The image on the left is immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using ml222276(EHHADH Antibody) at dilution 1/20, on the right is treated with fusion protein. (Original magnification: $\times 200$)

Western blotting

Predicted band size: 79 kDa

Positive control: Mouse liver tissue

Recommended dilution: 200-1000

Gel: 6%SDS-PAGE


Lysate: 40 μ g

Lane: Mouse liver tissue

Primary antibody: ml222276(EHHADH Antibody) at dilution 1/200

Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution

Exposure time: 1 minute



ELISA

Recommended dilution: 2000-5000

联系电话: 4008-898-798, 021-61725725

联系QQ: 2881505695, 2881505696

邮箱: mlbio_cn@yeah.net

网址: www.mlbio.cn