

订购热线: 4008-898-798

Anti-MYL12B antibody

| Cat. No. | ml222348 |
|--------------------|--|
| Package | 25 μl/100 μl/200 μl |
| Storage | -20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol |
| | |
| Product overview | |
| Description | Anti-MYL12B rabbit polyclonal antibody |
| Applications | ELISA, WB, IHC |
| Immunogen | Fusion protein of human MYL12B |
| Reactivity | Human, Mouse, Rat |
| Content | 0.6 mg/ml |
| Host species | Rabbit |
| lg class | Immunogen-specific rabbit IgG |
| Purification | Antigen affinity purification |
| | Antigen affinity purification |
| Target information | oroc |
| Symbol | MYL12B |
| Full name | myosin, light chain 12B, regulatory |
| | -,000 |
| | \smile |



Synonyms MLC-B; MRLC2

Swissprot 014950

Target Background

The activity of nonmuscle myosin II (see MYH9; MIM 160775) is regulated by phosphorylation of a regulatory light chain, such as MRLC2. This phosphorylation results in higher MgATPase activity and the assembly of myosin II filaments




Applications


Immunohistochemistry

Predicted cell location: Cytoplasm

Positive control: Human breast cancer

Recommended dilution: 50-200





The image on the left is immunohistochemistry of paraffin-embedded Human breast cancer tissue using ml222348(MYL12B Antibody) at dilution 1/25, on the right is treated with fusion protein. (Original magnification: ×200)

Western blotting


Predicted band size:20 kDa


Positive control:Mouse skeletal muscle, heart and bladder tissue

Predicted cell location: Cytoplasm

Positive control: Human prostate cancer

Recommended dilution: 50-200





The image on the left is immunohistochemistry of paraffin-embedded Human prostate cancer tissue using ml222348(MYL12B Antibody) at dilution 1/25, on the right is treated with fusion protein. (Original magnification: ×200)

Recommended dilution: 500-2000



Gel: 8%SDS-PAGE


Lysate: 40 µg

Lane 1-3: Mouse skeletal muscle, Mouse heart tissue, Mouse bladder tissue

Primary antibody: ml222348(MYL12B Antibody) at dilution 1/550

Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution

Exposure time: 20 seconds



ELISA

Recommended dilution: 2000-5000

联系电话: 4008-898-798, 021-61725725

联系QQ: 2881505695, 2881505696

邮箱: mlbio_cn@yeah.net

网址: www.mlbio.cn