

订购热线: 4008-898-798

Anti-RIOX2 antibody

Cat. No.	ml225067
Package	25 μl/100 μl/200 μl
Storage	-20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol
Product overview	
Description	Anti-RIOX2 rabbit polyclonal antibody
Applications	ELISA, WB, IHC
Immunogen	Fusion protein of human RIOX2
Reactivity	Human
Content	0.72 mg/ml
Host species	Rabbit
lg class	Immunogen-specific rabbit IgG
Purification	Antigen affinity purification
	AUCON
Target information	Antigen affinity purification
Symbol	RIOX2
Full name	ribosomal oxygenase 2
C	oou



Synonyms ROX; MDIG; MINA; NO52; JMJD10; MINA53

Swissprot Q8IUF8

Target Background

Oxygenase that can act as both a histone lysine demethylase and a ribosomal histidine hydroxylase. Is involved in the demethylation of trimethylated 'Lys-9' on histone H3 (H3K9me3), leading to an increase in ribosomal RNA expression. Also catalyzes the hydroxylation of 60S ribosomal protein L27a on 'His-39'. May play an important role in cell growth and survival. May be involved in ribosome biogenesis, most likely during the assembly process of pre-ribosomal particles.



订购热线: 4008-898-798


Applications

Immunohistochemistry

Predicted cell location: Nucleus

Positive control: Human tonsil

Recommended dilution: 25-100



The image on the left is immunohistochemistry of paraffin-embedded Human tonsil tissue using ml225067(RIOX2 Antibody) at dilution 1/20, on the right is treated with fusion protein. (Original magnification: ×200)

Western blotting

Predicted band size:53 kDa

Positive control:Jurkat cell lysate

Recommended dilution: 200-1000

Good elisakit producers



Gel: 8%SDS-PAGE


Lysate: 40 µg

Lane: Jurkat cell lysate

Primary antibody: ml225067(RIOX2 Antibody) at dilution 1/250

Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution

Exposure time: 45 seconds



ELISA

Recommended dilution: 5000-10000

联系电话: 4008-898-798, 021-61725725

联系QQ: 2881505695, 2881505696

邮箱: mlbio_cn@yeah.net

网址: www.mlbio.cn