

Anti-YWHAB antibody

Cat. No. ml260652

Package 25 μl/100 μl/200 μl

Storage -20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Product overview

Description Anti-YWHAB rabbit polyclonal antibody

Applications ELISA, WB, IHC

Immunogen Synthetic peptide of human YWHAB

Reactivity Human, Mouse, Rat

Content 0.4 mg/ml

Host species Rabbit

Ig class Immunogen-specific rabbit IgG

Purification Antigen affinity purification

Target information

Symbol YWHAB

Full name tyrosine 3-monooxygenase/tryptophan 5-monooxygenase activation protein, beta polypeptide



Synonyms HS1, GW128, YWHAA, KCIP-1

Swissprot P31946

Target Background

This gene encodes a protein belonging to the 14-3-3 family of proteins, members of which mediate signal transduction by binding to phosphoserine-containing proteins. This highly conserved protein family is found in both plants and mammals. The encoded protein has been shown to interact with RAF1 and CDC25 phosphatases, suggesting that it may play a role in linking mitogenic signaling and the cell cycle machinery. Two transcript variants, which encode the same protein, have been identified for this gene.




Applications


Immunohistochemistry

Predicted cell location: Cytoplasm, Nucleus

Positive control: Human lung cancer

Recommended dilution: 25-100





Good elisakii produceri

The image on the left is immunohistochemistry of paraffin-embedded Human lung cancer tissue using ml260652(YWHAB Antibody) at dilution 1/30, on the right is treated with synthetic peptide. (Original magnification: ×200)

Western blotting

Predicted band size:28 kDa


Positive control:Hela cells and mouse kidney tissue


Recommended dilution: 500-2000

Predicted cell location: Cytoplasm, Nucleus

Positive control: Human colon cancer

Recommended dilution: 25-100





The image on the left is immunohistochemistry of paraffin-embedded Human colon cancer tissue using ml260652(YWHAB Antibody) at dilution 1/30, on the right is treated with synthetic peptide. (Original magnification: ×200)



Gel: 12%SDS-PAGE


Lysate: 30 µg

Lane 1-2: Hela cells, mouse kidney tissue

Primary antibody: ml260652(YWHAB Antibody) at dilution 1/550

Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution

Exposure time: 10 seconds



ELISA

Recommended dilution: 1000-5000

联系电话: 4008-898-798, 021-61725725

联系QQ: 2881505695, 2881505696

邮箱: mlbio_cn@yeah.net

网址: www.mlbio.cn