

Anti-PAWR antibody

| | |
|-----------------|---|
| Cat. No. | ml260694 |
| Package | 25 µl/100 µl/200 µl |
| Storage | -20°C, pH7.4 PBS, 0.05% NaN ₃ , 40% Glycerol |

Product overview

| | |
|---------------------|--------------------------------------|
| Description | Anti-PAWR rabbit polyclonal antibody |
| Applications | ELISA, WB |
| Immunogen | Synthetic peptide of human PAWR |
| Reactivity | Human |
| Content | 0.3 mg/ml |
| Host species | Rabbit |
| Ig class | Immunogen-specific rabbit IgG |
| Purification | Antigen affinity purification |

Target information

| | |
|------------------|---------------------------------|
| Symbol | PAWR |
| Full name | PRKC, apoptosis, WT1, regulator |
| Synonyms | PAR4, Par-4 |
| Swissprot | Q96IZ0 |

Target Background

The tumor suppressor WT1 represses and activates transcription. The protein encoded by this gene is a WT1-interacting protein that itself functions as a transcriptional repressor. It contains a putative leucine zipper domain which interacts with the zinc finger DNA binding domain of WT1. This protein is specifically upregulated during apoptosis of prostate cells.

订购热线: 4008-898-798

Applications

Western blotting

Predicted band size: 37 kDa

Positive control: HeLa and A549 cells

Recommended dilution: 500-2000

Gel: 10% SDS-PAGE


Lysate: 30 µg

Lane 1-2: HeLa cells, A549 cells

Primary antibody: ml260694 (PAWR Antibody) at dilution 1/400

Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution

Exposure time: 10 seconds



ELISA

Recommended dilution: 2000-5000

联系电话: 4008-898-798, 021-61725725

联系QQ: 2881505695, 2881505696

邮箱: mlbio_cn@yeah.net

网址: www.mlbio.cn