

Anti-COX6A2 antibody

 Cat. No.
 ml261449

 Package
 25 μl/100 μl/200 μl

 Storage
 -20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol

| Product overview | |
|--------------------|--|
| Description | Anti-COX6A2 rabbit polyclonal antibody |
| Applications | ELISA, WB |
| Immunogen | Synthetic peptide of human COX6A2 |
| Reactivity | Human |
| Content | 0.2 mg/ml |
| Host species | Rabbit |
| lg class | Immunogen-specific rabbit IgG |
| Purification | Antigen affinity purification |
| | |
| Target information | |
| Symbol | COX6A2 |
| | |

COX6A2 cytochrome c oxidase subunit 6A2 COX6AH; COXVIAH Q02221

Target Background

Full name

Synonyms

Swissprot

Cytochrome c oxidase (COX), the terminal enzyme of the mitochondrial respiratory chain, catalyzes the electron transfer from reduced cytochrome c to oxygen. It is a heteromeric complex consisting of 3 catalytic subunits encoded by mitochondrial genes and multiple structural subunits encoded by nuclear genes. The mitochondrially-encoded subunits function in electron transfer, and the nuclear-encoded subunits may be involved in the regulation and assembly of the complex. This nuclear gene encodes polypeptide 2 (heart/muscle isoform) of subunit VIa, and polypeptide 2 is present only in striated muscles. Polypeptide 1 (liver isoform) of subunit VIa is encoded by a different gene, and is found in all non-muscle tissues. These two polypeptides share 66% amino acid sequence identity.



订购热线: 4008-898-798

Applications Western blotting Predicted band size:11 kDa Positive control:Human heart tissue lysate Recommended dilution: 200-1000

Gel: 12%SDS-PAGE Lysate: 40 µg Lane: Human heart tissue lysate Primary antibody: ml261449(COX6A2 Antibody) at dilution 1/200 Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution Exposure time: 1 second

ELISA

Recommended dilution:5000-10000

Wit produces 联系电话: 4008-898-798, 021-61725725

联系QQ: 2881505695, 2881505696

- 邮箱: mlbio_cn@yeah.net
- 网址: www.mlbio.cn