

Anti-CYP1A1 antibody

 Cat. No.
 ml262002

 Package
 25 μl/100 μl/200 μl

 Storage
 -20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Anti-CYP1A1 rabbit polyclonal antibody
ELISA, WB
Synthetic peptide of human CYP1A1
Human, Mouse, Rat
0.3 mg/ml
Rabbit
Immunogen-specific rabbit IgG
Antigen affinity purification
CYP1A1
cytochrome P450, family 1, subfamily A, polypeptide 1
AHH; AHRR; CP11; CYP1; P1-450; P450-C; P450DX
P04798

Target Background

This gene, CYP1A1, encodes a member of the cytochrome P450 superfamily of enzymes. The cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. This protein localizes to the endoplasmic reticulum and its expression is induced by some polycyclic aromatic hydrocarbons (PAHs), some of which are found in cigarette smoke. The enzyme's endogenous substrate is unknown; however, it is able to metabolize some PAHs to carcinogenic intermediates. The gene has been associated with lung cancer risk. A related family member, CYP1A2, is located approximately 25 kb away from CYP1A1 on chromosome 15.


res



订购热线: 4008-898-798

Applications Western blotting Predicted band size:58 kDa Positive control:Mouse lung tissue Recommended dilution: 200-1000

Gel: 6%SDS-PAGE Lysate: 40 µg Lane: Mouse lung tissue Primary antibody: ml262002(CYP1A1 Antibody) at dilution 1/220 Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution Exposure time: 30 seconds



ELISA

Recommended dilution:1000-2000

联系电话: 4008-898-798, 021-61725725

联系QQ: 2881505695, 2881505696

- 邮箱: mlbio_cn@yeah.net
- 网址: www.mlbio.cn