

订购热线: 4008-898-798

Anti-MED28 antibody

Cat. No. ml123403

Package 25 μ l/100 μ l/200 μ l

Storage -20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Product overview

Description Anti-MED28 rabbit polyclonal antibody

Applications ELISA, IHC

ImmunogenFull length fusion proteinReactivityHuman, Mouse, Rat

Content0.4 mg/mlHost speciesRabbit

Ig classImmunogen-specific rabbit IgGPurificationAntigen affinity purification

Target information

Symbol MED28

Full namemediator complex subunit 28SynonymsEG1; magicin; 1500003D12Rik

Swissprot Q9H204

Target Background

Component of the Mediator complex, a coactivator involved in the regulated transcription of nearly all RNA polymerase II-dependent genes. Mediator functions as a bridge to convey information from gene-specific regulatory proteins to the basal RNA polymerase II transcription machinery. Mediator is recruited to promoters by direct interactions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex with RNA polymerase II and the general transcription factors. May be part of a complex containing NF2/merlin that participates in cellular signaling to the actin cytoskeleton downstream of tyrosine kinase signaling pathways.




订购热线: 4008-898-798


Applications

Immunohistochemistry

Predicted cell location: Cytoplasm or Nucleus Positive control: Human thyroid cancer

Recommended dilution: 25-100





The image on the left is immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using ml123403(MED28 Antibody) at dilution 1/25, on the right is treated with fusion protein. (Original magnification: ×200)

ELISA

Recommended dilution: 2000-5000

联系电话: 4008-898-798, 021-61725725


联系QQ: 2881505695, 2881505696


邮箱: mlbio_cn@yeah.net 网址: www.mlbio.cn

Predicted cell location: Cytoplasm or Nucleus

Positive control: Human breast cancer

Recommended dilution: 25-100





The image on the left is immunohistochemistry of paraffin-embedded Human breast cancer tissue using ml123403(MED28 Antibody) at dilution 1/25, on the right is treated with fusion protein. (Original

magnification: ×200)