

Anti-MIF antibody

| | |
|-----------------|---|
| Cat. No. | ml263076 |
| Package | 25 µl/100 µl/200 µl |
| Storage | -20°C, pH7.4 PBS, 0.05% NaN ₃ , 40% Glycerol |

Product overview

| | |
|---------------------|-------------------------------------|
| Description | Anti-MIF rabbit polyclonal antibody |
| Applications | ELISA, WB |
| Immunogen | Synthetic peptide of human MIF |
| Reactivity | Human, Mouse, Rat |
| Content | 0.4 mg/ml |
| Host species | Rabbit |
| Ig class | Immunogen-specific rabbit IgG |
| Purification | Antigen affinity purification |

Target information

| | |
|------------------|--|
| Symbol | MIF |
| Full name | macrophage migration inhibitory factor (glycosylation-inhibiting factor) |
| Synonyms | GIF; GLIF; MMIF |
| Swissprot | P14174 |

Target Background

This gene encodes a lymphokine involved in cell-mediated immunity, immunoregulation, and inflammation. It plays a role in the regulation of macrophage function in host defense through the suppression of anti-inflammatory effects of glucocorticoids. This lymphokine and the JAB1 protein form a complex in the cytosol near the peripheral plasma membrane, which may indicate an additional role in integrin signaling pathways.

订购热线: 4008-898-798

Applications

Western blotting

Predicted band size:12 kDa

Positive control:HT-29, HeLa and Jurkat cell, Rat brain tissue lysates

Recommended dilution: 500-2000

Gel: 12%SDS-PAGE


Lysate: 40 μ g

Lane 1-4: HT-29, HeLa and Jurkat cell, Rat brain tissue lysates

Primary antibody: ml263076(MIF Antibody) at dilution 1/200

Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution

Exposure time: 50 seconds



ELISA

Recommended dilution:2000-5000

联系电话: 4008-898-798, 021-61725725

联系QQ: 2881505695, 2881505696

邮箱: mlbio_cn@yeah.net

网址: www.mlbio.cn