

兔抗 AADACL2 多克隆抗体

中文名称: 兔抗 AADACL2 多克隆抗体

英文名称: Anti-AADACL2 rabbit polyclonal antibody

储 存: 冷冻(-20℃)

抗 原: AADACL2

宿 主: Rabbit

相关类别: 一抗

反应种属: Human

标记物: Unconjugate

克隆类型: rabbit polyclonal

技术规格

Background:

AADAC is a 45 kDa membrane-associated hydrolase highly expressed in the human liver that has been im plicated in the hydrolysis of arylacetamide xenobiotics including flutamide and phenacetin.556–560AADAC is highly homologous to several related lipases, AADACL 1, 2, and 4, the latter two of which remain completel y unannotated. Overexpression of AADAC in a rat he patoma cell line produces changes in cellular TG level s, apoplipoprotein B secretion, and fatty acid oxidatio n, but the mechanism for these effects remains uncle ar.561 Candidate endogenous substrates, selective inh ibitors, or knockout mice have not, to our knowledge, been described for AADAC or the related enzymes



	AADACL2 and AADACL4.
Applications:	ELISA, IHC
Name of antibody:	AADACL2
Immunogen:	Fusion protein of human AADACL2
Full name:	arylacetamide deacetylase-like 2
SwissProt:	Q6P093
ELISA Recommended dilution:	2000-5000
IHC positive control:	Human cervical cancer and human thyroid cancer
IHC Recommend dilution:	25-100

