

## 兔抗 ACSBG2 多克隆抗体

中文名称：兔抗 ACSBG2 多克隆抗体

英文名称：Anti-ACSBG2 rabbit polyclonal antibody

别名：BGR, BRGL, PRDNY3, PRD-NY3

相关类别：一抗

储存：冷冻（-20℃）

抗原：ACSBG2

宿主：Rabbit

反应种属：Human

标记物：Unconjugate

克隆类型：rabbit polyclonal

### 技术规格

<b>Background:</b>	Long-chain-fatty-acid—CoA ligase ACSBG2 is an enzyme that in humans is encoded by the ACSBG2 gene. Mediates activation of long-chain fatty acids for both synthesis of cellular lipids, and degradation via beta-oxidation. Able to activate long-chain fatty acids. Also able to activate very long-chain fatty acids, however, the relevance of such activity is unclear in vivo. Has increased ability to activate oleic and linoleic acid. May play a role in spermatogenesis.
<b>Applications:</b>	ELISA, IHC
<b>Name of antibody:</b>	ACSBG2

<b>Immunogen:</b>	Fusion protein of human ACSBG2
<b>Full name:</b>	acyl-CoA synthetase bubblegum family member 2
<b>Synonyms :</b>	BGR, BRGL, PRTDNY3, PRTD-NY3
<b>SwissProt:</b>	Q5FVE4
<b>ELISA Recommended dilution:</b>	500-5000
<b>IHC positive control:</b>	Human stomach cancer
<b>IHC Recommend dilution:</b>	50-100

