

兔抗 BAG1 多克隆抗体

- 中文名称: 兔抗 BAG1 多克隆抗体
- 英文名称: Anti-BAG1 rabbit polyclonal antibody
- 别名: HAP; BAG-1; RAP46
- 储存: 冷冻(-20℃)
- 抗原: BAG1
- 宿 主: Rabbit
- 相关类别: 一抗
- 反应种属: Human
- 标记物: Unconjugate
- 克隆类型: rabbit polyclonal

技术规格

Background:	The oncogene BCL2 is a membrane protein that block s a step in a pathway leading to apoptosis or progra mmed cell death. The protein encoded by this gene b inds to BCL2 and is referred to as BCL2-associated ath anogene. It enhances the anti-apoptotic effects of BCL 2 and represents a link between growth factor recepto rs and anti-apoptotic mechanisms. Multiple protein iso forms are encoded by this mRNA through the use of a non-AUG (CUG) initiation codon, and three alternativ e downstream AUG initiation codons. A related pseud
	ogene has been defined on chromosome X.



Applications:	ELISA, WB, IHC
Name of antibody:	BAG1
Immunogen:	Fusion protein of human BAG1
Full name:	BCL2-associated athanogene
Synonyms :	HAP; BAG-1; RAP46
SwissProt:	Q99933
ELISA Recommended dilution:	2000-5000
IHC positive control:	Human cervical cancer and human thyroid cancer
IHC Recommend dilution:	25-100
WB Predicted band size:	39 kDa
WB Positive control:	Hela and Raji cells
WB Recommended dilution:	500-2000





