

## 兔抗 CXCL14 多克隆抗体

中文名称：兔抗 CXCL14 多克隆抗体

英文名称：Anti-CXCL14 rabbit polyclonal antibody

别 名：KEC; KS1; BMAC; BRAK; NJAC; MIP2G; MIP-2g; SCYB14

抗 原：CXCL14

储 存：冷冻（-20℃）

宿 主：Rabbit

相关类别：一抗

反应种属：Human

标记物：Unconjugate

克隆类型：rabbit polyclonal

### 技术规格

#### Background:

This gene belongs to the cytokine gene family which encode secreted proteins involved in immunoregulatory and inflammatory processes. The protein encoded by this gene is structurally related to the CXC (Cys-X-Cys) subfamily of cytokines. Members of this subfamily are characterized by two cysteines separated by a single amino acid. This cytokine displays chemotactic activity for monocytes but not for lymphocytes, dendritic cells, neutrophils or macrophages. It has been implicated that this cytokine is involved in the homeostasis of monocyte-derived macrophages rather than

|                                    |   |
|------------------------------------|---|
|                                    | in inflammation.                                  |
| <b>Applications:</b>               | ELISA, IHC  |
| <b>Name of antibody:</b>           | CXCL14  |
| <b>Immunogen:</b>                  | Fusion protein of human CXCL14                    |
| <b>Full name:</b>                  | chemokine (C-X-C motif) ligand 14                 |
| <b>Synonyms :</b>                  | KEC; KS1; BMAC; BRAK; NJAC; MIP2G; MIP-2g; SCYB14 |
| <b>SwissProt:</b>                  | O95715  |
| <b>ELISA Recommended dilution:</b> | 2000-5000   |
| <b>IHC positive control:</b>       | Human thyroid cancer                              |
| <b>IHC Recommend dilution:</b>     | 50-200  |

