

RNF146 抗原(重组蛋白)

中文名称: RNF146 抗原(重组蛋白)

- 英文名称: RNF146 Antigen (Recombinant Protein)
- 别名: ring finger protein 146
- 储存: 冷冻 (-20℃)
- 相关类别: 抗原

概述

Fusion protein corresponding to a region derived from 160-359 amino acids of human RNF146

技术规格

Full name:	ring finger protein 146
Swissprot:	Q9NTX7
Gene Accession:	BC008235
Purity:	>85%, as determined by Coomassie blue stained SDS-PAGE
Expression system:	Escherichia coli
Tags:	His tag C-Terminus, GST tag N-Terminus
Background:	E3 ubiquitin-protein ligase that specifically binds poly-ADP-ribosylated (PARsylated) proteins and mediates their ubiquitination and subsequen t degradation. May regulate many important biological processes, such as cell survival and DNA damage response. Acts as an activator of the Wnt signaling pathway by mediating the ubiquitination of PARsylated AXIN1 and AXIN2, 2 key components of the beta-catenin destruction complex. Acts in cooperation with tankyrase proteins (TNKS and TNKS 2), which mediate PARsylation of target proteins AXIN1, AXIN2, BLZF1, CASC3, TNKS and TNKS2. Recognizes and binds tankyrase-dependent



PARsylated proteins via its WWE domain and mediates their ubiquitina tion, leading to their degradation. Different ubiquitin linkage types hav e been observed: TNKS2 undergoes ubiquitination at 'Lys-48' and 'Lys-63', while AXIN1 is only ubiquitinated at 'Lys-48'. May regulate TNKS and TNKS2 subcellular location, preventing aggregation at a centroso mal location. Neuroprotective protein. Protects the brain against N-me thyl-D-aspartate (NMDA) receptor-mediated glutamate excitotoxicity an d ischemia, by interfering with PAR-induced cell death, called parthana tos. Prevents nuclear translocation of AIFM1 in a PAR-binding depend ent manner. Does not affect PARP1 activation (By similarity). Protects against cell death induced by DNA damaging agents, such as N-meth yl-N-nitro-N-nitrosoguanidine (MNNG) and rescues cells from G1 arres t. Promotes cell survival after gamma-irradiation. Facilitates DNA repair