

RPLP0 抗原（重组蛋白）

中文名称：RPLP0 抗原（重组蛋白）

英文名称：RPLP0 Antigen (Recombinant Protein)

别名：P0; LP0; L10E; RPP0; PRLP0

储存：冷冻（-20℃）

相关类别：抗原

概述

Full length fusion protein

技术规格

Full name:	Ribosomal protein, large, P0
Synonyms:	P0; LP0; L10E; RPP0; PRLP0
Swissprot:	P05388
Gene Accession:	BC000087
Purity:	>85%, as determined by Coomassie blue stained SDS-PAGE
Expression system:	Escherichia coli
Tags:	His tag C-Terminus, GST tag N-Terminus
Background:	Ribosomes, the organelles that catalyze protein synthesis, consist of a small 40S subunit and a large 60S subunit. Together these subunits are composed of 4 RNA species and approximately 80 structurally distinct proteins. This gene encodes a ribosomal protein that is a component of the 60S subunit. The protein, which is the functional equivalent of the E. coli L10 ribosomal protein, belongs to the L10P family of ribosomal proteins. It is a neutral phosphoprotein with a C-terminal e

nd that is nearly identical to the C-terminal ends of the acidic ribosomal phosphoproteins P1 and P2. The P0 protein can interact with P1 and P2 to form a pentameric complex consisting of P1 and P2 dimers, and a P0 monomer. The protein is located in the cytoplasm. Transcript variants derived from alternative splicing exist; they encode the same protein. As is typical for genes encoding ribosomal proteins, there are multiple processed pseudogenes of this gene dispersed throughout the genome.