

兔抗 CBL (phospho-Tyr700) 多克隆抗体

中文名称：兔抗 CBL (phospho-Tyr700) 多克隆抗体

英文名称：Anti-CBL (phospho-Tyr700) rabbit polyclonal antibody

别名：CBL2; NSLL; C-CBL; RNF55; FRA11B

相关类别：一抗

储存：冷冻（-20℃）避光

宿主：Rabbit

抗原：CBL (phospho-Tyr700)

反应种属：Human, Mouse, Rat

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

Background:	Participates in signal transduction in hematopoietic cells. Adapter protein that functions as a negative regulator of many signaling pathways that start from receptors at the cell surface. Acts as an E3 ubiquitin-protein ligase, which accepts ubiquitin from specific E2 ubiquitin-conjugating enzymes, and then transfers it to substrates promoting their degradation by the proteasome. Recognizes activated receptor tyrosine kinases, including PDGFA, EGF and CSF1, and terminates signaling.
Applications:	WB, IHC, IF
Name of antibody:	CBL (phospho-Tyr700)

Immunogen:	Synthetic peptide of human CBL (phospho-Tyr700)
Full name:	Cbl proto-oncogene, E3 ubiquitin protein ligase (phospho-Tyr700)
Synonyms :	CBL2; NSLL; C-CBL; RNF55; FRA11B
SwissProt:	P22681
IHC positive control:	Human breast carcinoma
IHC Recommend dilution:	50-100
WB Predicted band size:	120 kDa
WB Positive control:	HT29 cells treated with UV
WB Recommended dilution:	500-1000
IF Positive control:	Hela cells
IF Recommended dilution	100-200





