

兔抗 CD3G 多克隆抗体

- 中文名称：兔抗 CD3G 多克隆抗体
- 英文名称：Anti-CD3G rabbit polyclonal antibody
- 别名：CD3g molecule; T3G; IMD17; CD3-GAMMA
- 相关类别：一抗
- 储存：冷冻（-20℃）
- 宿主：Rabbit
- 抗原：CD3G
- 反应种属：Human, Mouse
- 标记物：Unconjugate
- 克隆类型：rabbit polyclonal

技术规格

Background:	The protein encoded by this gene is the CD3-gamma polypeptide, which together with CD3-epsilon, -delta and -zeta, and the T-cell receptor alpha/beta and gamma/delta heterodimers, forms the T-cell receptor-CD3 complex. This complex plays an important role in coupling antigen recognition to several intracellular signal-transduction pathways. The genes encoding the epsilon, gamma and delta polypeptides are located in the same cluster on chromosome 11. Defects in this gene are associated with T cell immunodeficiency.
Applications:	ELISA, WB

Name of antibody:	CD3G
Immunogen:	Fusion protein of human CD3G
Full name:	CD3g molecule
Synonyms:	T3G; IMD17; CD3-GAMMA
SwissProt:	P09693
ELISA Recommended dilution:	5000-10000
WB Predicted band size:	20 kDa
WB Positive control:	Mouse thymus tissue lysate
WB Recommended dilution:	200-1000

