

兔抗 CD70 多克隆抗体

中文名称：兔抗 CD70 多克隆抗体

英文名称：Anti-CD70 rabbit polyclonal antibody

别名：CD70; CD27L; CD27LG; TNFSF7; CD70 antigen; Tumor necrosis factor ligand superfamily member 7

相关类别：一抗

储存：冷冻（-20℃）避光

宿主：Rabbit

抗原：CD70

反应种属：Human, Mouse, Rat

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

Background:	The protein encoded by this gene is a cytokine that belongs to the tumor necrosis factor (TNF) ligand family. This cytokine is a ligand for TNFRSF27/CD27. It is a surface antigen on activated, but not on resting, T and B lymphocytes. It induces proliferation of costimulated T cells, enhances the generation of cytolytic T cells, and contributes to T cell activation. This cytokine is also reported to play a role in regulating B-cell activation, cytotoxic function of natural killer cells, and immunoglobulin synthesis.
Applications:	WB

Name of antibody:	CD70
Immunogen:	Fusion protein of human CD70
Full name:	CD70 molecule
Synonyms :	CD70; CD27L; CD27LG; TNFSF7; CD70 antigen; Tumor necrosis factor ligand superfamily member 7
SwissProt:	P32970
WB Predicted band size:	21 kDa
WB Positive control:	NCI-H460 cells
WB Recommended dilution:	500-2000



