



兔抗 ELOC 多克隆抗体

中文名称：兔抗 ELOC 多克隆抗体

英文名称：Anti-ELOC rabbit polyclonal antibody

别 名：elongin C; SIII; TCEB1

相关类别：一抗

储 存：冷冻 (-20°C)

宿 主：Rabbit

抗 原：ELOC

反应种属：Human, Mouse, Rat

标记物：Unconjugate


克隆类型：rabbit polyclonal

技术规格

Background:

This gene encodes the protein elongin C, which is a subunit of the transcription factor B (SIII) complex. The SII I complex is composed of elongins A/A2, B and C. It activates elongation by RNA polymerase II by suppressing transient pausing of the polymerase at many sites within transcription units. Elongin A functions as the transcriptionally active component of the SIII complex, whereas elongins B and C are regulatory subunits. Elongin A2 is specifically expressed in the testis, and capable of forming a stable complex with elongins B and C. The von Hippel-Lindau tumor suppressor protein binds to elongins

| | |
|------------------------------------|--|
| | B and C, and thereby inhibits transcription elongation. Multiple alternatively spliced transcript variants encoding two distinct isoforms have been identified. |
| Applications: | ELISA, IHC |
| Name of antibody: | ELOC |
| Immunogen: | Fusion protein of human ELOC |
| Full name: | elongin C |
| Synonyms: | SIII; TCEB1 |
| SwissProt: | Q15369 |
| ELISA Recommended dilution: | 5000-10000 |
| IHC positive control: | Human breast cancer Human ovarian cancer |
| IHC Recommend dilution: | 50-300 |



mlbio 海聯生物
Good elisakit producers

