

## PPM1D 抗原(重组蛋白)

中文名称: PPM1D 抗原(重组蛋白)

英文名称: PPM1D Antigen (Recombinant Protein)

别 名: protein phosphatase, Mg2+/Mn2+ dependent 1D; WIP1; PP2C-DELTA

储 存: 冷冻(-20℃)

相关类别: 抗原

概 述:

Fusion protein corresponding to a region derived from 459-605 amino acids of human PPM1D

## 技术规格:

| Full name:         | protein phosphatase, Mg2+/Mn2+ dependent 1D  |
|--------------------|--|
| Synonyms:          | WIP1; PP2C-DELTA   |
| Swissprot:         | O15297   |
| Gene Accession:    | BC033893   |
| Purity:            | >85%, as determined by Coomassie blue stained SDS-PAGE   |
| Expression system: | Escherichia coli   |
| Tags:              | His tag C-Terminus, GST tag N-Terminus   |
| Background:        | The protein encoded by this gene is a member of the PP2C family of Ser/Thr protein phosphatases. PP2C family members are known to be negative regulators of cell stress response pathways. The expression of this gene is induced in a p53-dependent manner in response to vario us environmental stresses. While being induced by tumor suppressor p rotein TP53/p53, this phosphatase negatively regulates the activity of p38 MAP kinase, MAPK/p38, through which it reduces the phosphoryl |



ation of p53, and in turn suppresses p53-mediated transcription and a poptosis. This phosphatase thus mediates a feedback regulation of p3 8-p53 signaling that contributes to growth inhibition and the suppress ion of stress induced apoptosis. This gene is located in a chromosoma I region known to be amplified in breast cancer. The amplification of this gene has been detected in both breast cancer cell line and prima ry breast tumors, which suggests a role of this gene in cancer development.