

兔抗 FSD1L 多克隆抗体

中文名称：兔抗 FSD1L 多克隆抗体

英文名称：Anti-FSD1L rabbit polyclonal antibody

别 名：MIR1; CCDC10; FSD1CL; FSD1NL; CSDUFD1

相关类别：一抗

储 存：冷冻（-20℃）

宿 主：Rabbit

抗 原：FSD1L

反应种属：Human, Mouse

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

Background:

FSD1L (fibronectin type III and SPRY domain containing 1-like), also known as CCDC10 (coiled-coil domain-containing protein 10), CSDUFD1, MIR1 or FSD1CL, is a 530 amino acid protein containing one B30.2/SPRY domain, one COS domain, and a fibronectin type-III domain. Existing as three alternatively spliced isoforms, FSD1L is expressed primarily in brain, with lower levels of expression found in thymus, pituitary and testis. FSD1L may function in microtubule binding during interphase and is encoded by a gene th

	at maps to human chromosome 9q31.2. Chromosome 9 consists of about 145 million bases and comprises approximately 4% of the human genome and encodes nearly 900 genes. Considered to play a role in gender determination, deletion of the distal portion of 9p can lead to development of male to female sex reversal, the phenotype of a female with a male X,Y genotype.
Applications:	ELISA, IHC
Name of antibody:	FSD1L
Immunogen:	Fusion protein of human FSD1L
Full name:	fibronectin type III and SPRY domain containing 1-like
Synonyms :	MIR1; CCDC10; FSD1CL; FSD1NL; CSDUFD1
SwissProt:	Q9BXM9
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human esophagus cancer and human thyroid cancer
IHC Recommend dilution:	25-100

