

兔抗 GLYCTK 多克隆抗体

中文名称: 兔抗 GLYCTK 多克隆抗体

英文名称: Anti-GLYCTK rabbit polyclonal antibody

别 名: HBEBP2; HBEBP4; HBeAgBP4A

相关类别: 一抗

储 存: 冷冻(-20℃) 避光

抗 原: GLYCTK

宿 主: Rabbit

反应种属: Human Mouse Rat

标记物: Unconjugate

克隆类型: Unconjugate

技术规格

Background:	This locus encodes a member of the glycerate kinas e type-2 family. The encoded enzyme catalyzes the phosphorylation of (R)-glycerate and may be involve d in serine degradation and fructose metabolism. D ecreased activity of the encoded enzyme may be as sociated with the disease D-glyceric aciduria. Alterna tively spliced transcript variants have been described .
Applications:	WB, IHC, IF
Name of antibody:	GLYCTK
Immunogen:	Synthesized peptide derived from internal of human



	GLCTK.
Full name:	glycerate kinase
Synonyms:	HBEBP2; HBEBP4; HBeAgBP4A
SwissProt:	Q8IVS8
IHC positive control:	Human brain tissue
IHC Recommend dilution:	50-100
WB Predicted band size:	55 kDa
WB Positive control:	NIH/3T3 cells lysate
WB Recommended dilution:	500-3000
IF positive control:	A549 cells
IF Recommend dilution:	100-500





