

兔抗 MAP3K5(Ab-966)多克隆抗体

- 中文名称: 兔抗 MAP3K5(Ab-966)多克隆抗体
- 英文名称: Anti-MAP3K5(Ab-966) rabbit polyclonal antibody
- 别 名: ASK1; MEKK5; MAPKKK5
- 相关类别: 一抗
- 储 存: 冷冻 (-20℃)
- 抗 原: MAP3K5(Ab-966)
- 宿 主: Rabbit
- 反应种属: Human, Mouse, Rat
- 标记物: Unconjugate
- 克隆类型: rabbit polyclonal

技术规格

IF Recommended dilution	100-200
IF Positive control:	Hela cells
WB Recommended dilution:	500-1000
WB Positive control:	293 and Hela cell lysates
WB Predicted band size:	155 kDa
IHC Recommend dilution:	50-100
IHC positive control:	Human breast carcinoma
SwissProt:	Q99683
Synonyms:	ASK1; MEKK5; MAPKKK5
Full name:	mitogen-activated protein kinase kinase kinase 5



Immunogen:	Peptide sequence around aa.964-968 (S-I-S-L-P) derived fro m Human MAP3K5
Name of antibody:	MAP3K5(Ab-966)
Applications:	WB, IHC, IF
Background:	Mitogen-activated protein kinase (MAPK) signaling cascades include MAPK or extracellular signal-regulated kinase (ERK), MAPK kinase (MKK or MEK), and MAPK kinase kinase (MAP KKK or MEKK). MAPKK kinase/MEKK phosphorylates and act ivates its downstream protein kinase, MAPK kinase/MEK, wh ich in turn activates MAPK. The kinases of these signaling c ascades are highly conserved, and homologs exist in yeast, Drosophila, and mammalian cells. MAPKKK5 contains 1,374 amino acids with all 11 kinase subdomains. Northern blot a nalysis shows that MAPKKK5 transcript is abundantly expres sed in human heart and pancreas. The MAPKKK5 protein p hosphorylates and activates MKK4 (aliases SERK1, MAPKK4) in vitro, and activates c-Jun N-terminal kinase (JNK)/stress-a ctivated protein kinase (SAPK) during transient expression in COS and 293 cells; MAPKKK5 does not activate MAPK/ERK.









