



# 兔抗 HDAC2 (Phospho-Ser394)多克隆抗体

中文名称： 兔抗 HDAC2 (Phospho-Ser394)多克隆抗体

英文名称： Anti-HDAC2 (Phospho-Ser394) rabbit polyclonal antibody

别 名： HD2; RPD3; YAF1

相关类别： 一抗

储 存： 冷冻 (-20°C) 避光

宿 主： Rabbit

抗 原： HDAC2 (Phospho-Ser394)

反应种属： Human, Mouse, Rat


标记物： Unconjugate

克隆类型： rabbit polyclonal

## 技术规格

|                          |  |
|--------------------------|--|
| <b>Background:</b>       | Responsible for the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4). Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events. Histone deacetylases act via the formation of large multiprotein complexes |
| <b>Applications:</b>     | WB, IHC, IF  |
| <b>Name of antibody:</b> | HDAC2 (Phospho-Ser394)   |
| <b>Immunogen:</b>        | Synthetic peptide of human HDAC2 (Phospho-Ser394)  |
| <b>Full name:</b>        | histone deacetylase 2 (Phospho-Ser394)   |
| <b>Synonyms :</b>        | HD2; RPD3; YAF1  |

|                                 |   |
|---------------------------------|---|
| <b>SwissProt:</b>               | Q92769                                  |
| <b>IHC positive control:</b>    | Human breast carcinoma                  |
| <b>IHC Recommend dilution:</b>  | 50-100                                  |
| <b>WB Predicted band size:</b>  | 55 kDa                                  |
| <b>WB Positive control:</b>     | 293 cells untreated or treated with EGF |
| <b>WB Recommended dilution:</b> | 500-1000                                |
| <b>IF Positive control:</b>     | Hela cells                              |
| <b>IF Recommended dilution</b>  | 100-200                                 |



**mlbio** 海聯生物  
Good elisakit producers

